
v .	Dear Parents,
1	Şo that I am able to provide your child with a severe alleray the best care possible. I am asking you to
·	
<u></u>	
i A	
<u>.</u>	· t
·	
) T	
t _{ie} .	

376-4510 or Jamie.miller@dvusd.org at any time with questions or concerns.

Your assistance is invaluable to me and I greatly appreciate it. Please feel free to contact me at 623-





Ť	en.					
	1					1
					\ <u>\</u>	
				4	_	
I.e.						
, ==						
7 4	-				<u> </u>	<u> </u>
ni çe						
					e	
<u> </u>						
			-			
			¥		_	'
	7.					
T						
in a second						
			-			-
		•				
<u> </u>						
· · · · · · · · · · · · · · · · · · ·		**				
			.			•
2	P					
T.			I			
			•			
î .						
, 						
			v			
*						
į	-					
<u> </u>						
-						
1						
:						
- - -		,				
*(
						,
3						

The prevalence of anaphylaxis is increasing. Epinephrine is the only recognized first-line treatment for anaphylaxis and must be given promptly. Epinephrine is underutilized in anaphylaxis treatment. All other drugs have a delayed onset of action. EPIPEN (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS: Remove the Eninen Auto-Injector from the plastic carrying case. Remove the BLUE safety release cap.